

Wheat Crop Development in Central Punjab (Faisalabad, 2022–23)



By

Ali Imran

(Meteorologist)

Regional Agrometeorological Centre

Pakistan Meteorological Department

Ayub Agriculture Research Institute

Faisalabad, Pakistan

Contents

ABSTRACT	6
CHAPTER 1: INTRODUCTION -----	7
1.1 Geographical Description and Climate of Pakistan and Central Punjab (Study Area)	7
1.2 Scope of the Study	8
1.3 Objective of the Study.....	9
1.4 Review of Agriculture Production in Pakistan.....	9
1.5 Wheat Production in Pakistan	9
CHAPTER 2: MATERIALS AND METHOD -----	10
2.1 Phenology	10
2.1.1 Phenological Observations	11
2.2 Methodology	12
CHAPTER 3: RESULTS AND DISCUSSION -----	13
3.1 Rainfall and Wheat Crop Growth	14
3.2 Irrigation during Crop Growth	16
3.3 Air Temperature and Wheat Crop Growth	16
3.4 Soil Moisture Observations during Crop Growth	18
3.5 Soil Temperature and Crop Growth	20
3.6 Heat Units Consumption and Crop Cycle.....	21
3.7 Relative Humidity (%)	23
3.8 Reference Crop Evapotranspiration, ETo (mm/day)	24
3.9 Wind and Crop Growth.....	24
3.10 Crop Water Requirement.....	25
3.11 Agro Meteorological Summary of Crop Cycle	28
CHAPTER 4: CONCLUSION AND RECOMMENDATIONS -----	30
4.1 Conclusions	31
4.2 Recommendations	31
REFERENCES	33

List of Figures

		Page No.
Figure 1.1	Mean Annual Rainfall (mm) of Punjab [Normal (1981-2010)]	8
Figure 1.2	Normal (1981-2010) Mean Monthly Rainfall (mm) of Faisalabad during Rabi Season	8
Figure 1.3	Normal (1981-2010) Mean daily Maximum and Minimum Temperature (°C) of Faisalabad during Rabi Season	8
Figure 3.1	Comparison of Monthly Observed Precipitation with Normal (1981-2010) at Faisalabad	14
Figure 3.2	Mean Monthly Minimum Temperature (°C) of Faisalabad.....	16
Figure 3.3	Mean Monthly Maximum Temperature (°C) of Faisalabad.....	16
Figure 3.4	Mean Monthly Temperature (°C) during 2022-23.....	17
Figure 3.5	Soil moisture chrono Isopleths for Rabi Crop at Faisalabad for the year 2022-23.....	19
Figure 3.6	Soil moisture at different depths during wheat crop 2022-23.....	19
Figure 3.7	Moisture at deep soils (90cm) during wheat crop 2022-23.....	20
Figure 3.8	Soil temperature during wheat crop 2022-23.....	21
Figure 3.9	Heat units during crop life 2022-23.....	22
Figure 3.10	March of Crop Coefficient (Kc) for normal duration of Wheat growing season (Emergence to Wax- Maturity).	26
Figure 3.11	Crop Water Requirement (CWR) During Crop Life.....	27

List of Tables

		Page No.
Table 2.1	Observed Meteorological Parameters	10
Table 2.2	Phenological Stages of Wheat Crop	11
Table 3.1	Brief Summary of the Wheat Crop	13
Table 3.2	Daily rainfall history of the crop life 2022-23.....	15
Table 3.3	Mean Monthly Temperature during Rabi Season 2022-23.....	17
Table 3.4	Heat Units Consumed by the Crop during Different Phenological Phases	23
Table 3.5	Summary of some Meteorological Parameters during Rabi Season 2022-23	24

Abstract

This study was conducted at RAMC (Regional Agrometeorological Center) Faisalabad to investigate the impact of weather conditions on the growth and development of wheat crop. Impact of variations in the meteorological parameters on different phenological phases and hence on final yield of wheat crop was analyzed. For this purpose, both meteorological and phenological data along with soil moisture data was collected from the meteorological observatory and agricultural field of physiology section of Ayub Agriculture Research Institute, Faisalabad. Besides this record, other necessary features like sowing time, fertilizer, weeds removing operations and irrigation schedule are also included in the current report for the Rabi season 2022-23. The wheat crop variety Arooj-2022 sown in the field under observation used 1695 heat units in 143 days during its life cycle from emergence to full maturity. The water requirement of crop was fulfilled by flood irrigation method. Crop was sown at the proper time. Air temperature remained above normal values during the most of the crop life. Rainfall received remained well below normal values during the crop season.

Chapter 1

INTRODUCTION

Wheat is the major food crop as well as one of the main agricultural products of Pakistan. This study is based upon field observations of wheat crop at Regional Agrometeorological Centre, Faisalabad cultivated in the experimental field of Ayub Agriculture research Institute, Faisalabad. Land at the study site consists of agriculture fields rotated among wheat, maize, rice, cotton and other seasonal crops. In experimental field wheat crop variety Arooj-2022 was cultivated. This study will help in understanding the effect of different climatic parameters on the growth and yield of wheat crop in central Punjab region.

1.1 Geographical Description and Climate of Central Punjab (Study Area)

Pakistan has a variable climate, ranging from arid (33-254mm annual rainfall) in the south to semi-arid (254-508mm annual rainfall), sub-humid (508-1016mm annual rainfall) and humid (1016-2032mm annual rainfall) in the north. The Indus River that originates from the north along with its tributaries irrigates the great plains of the country including Central Punjab. Chaudhry and Rasul found that about 2/3 of the total agriculture area lies in the arid climate. [1]

The study has been conducted for “Faisalabad District” in Central Punjab with longitude 73.06°E, latitude 31.26°N and elevation of 184.5m from mean sea level. The climate of central and southern Punjab possesses the dry semi-arid agro-climatic characteristics (Figure– 1.1) but well managed canal irrigation system has placed it among the highly productive agriculture zones. Summer monsoon produces more rainfall and winter has a little contribution. Day time temperature reaches above 40°C during summer from April to September except some occasional relief from monsoon rains and decreases the evaporation demand of the atmosphere. The winter season starts from November and continues till March. December, January and February are the coldest months. In winter, night time temperature drops below 0°C [2]. More detail about the climate of Faisalabad during Rabi season is shown in the figures (1.2& 1.3). The highest amount of rainfall occurs during Rabi season in the months of March followed by April and February respectively. Day time mean maximum and night time mean minimum temperature gradually decrease from November to January and then start rising [3].




Figure 1.1: Mean Annual Rainfall (mm) of Punjab
[Normal (1981-2010)]




Figure 1.2: Normal (1981-2010) Mean Monthly Rainfall (mm) of Faisalabad during Rabi Season




Figure 1.3: Normal (1981-2010) Mean daily Maximum and Minimum Temperature (°C) of Faisalabad during Rabi Season

1.2 Scope of the Study

Pakistan experiences a more variable and unpredictable behavior of weather systems as compared to other countries of the region. This type of weather patterns results in the variations in the amount of available water both for irrigated and rain fed agricultural lands, which consequently results in the fluctuations in annual yield of wheat. Sometime heavy rains along with persistent cloudy conditions trigger the viral or pest attack on wheat crop and also cause rapid growth of weeds in the fields. It also prolongs the crop period, which causes delay in sowing of next Kharif crop. Rain just after sowing causes decrease in the number of germinated wheat seeds and also at the time of harvesting/threshing badly damage the final yield. Abnormal rise in day time temperature or late sowing raises crop water requirement at a particular

phase. This could also cause early completion of phase and early maturity of grains. Due to which shriveled grain is obtained. Therefore, in this study the impacts of variations of meteorological parameters along with variations in soil temperature and moisture are analyzed to understand crop growth and development throughout the crop life and their impact on final yield of the crop.

1.3 Objective of the Study

- To analyze and study the impact of various meteorological parameters on crop growth and development.
- To investigate the water satisfaction sensitivity of crop in Faisalabad region.
- To develop the relationship between weather parameters, crops life cycle and yield obtained.
- To determine the onset of pests and diseases related to weather elements.
- To get a step forward for formulation of yield estimation.

1.4 Review of Agriculture Production in Pakistan

Pakistan is an agrarian country whose population and economy directly or indirectly (70% directly and 16% indirectly) depends upon agriculture. Agriculture is the mainstay of Pakistan's economy. It accounts for 21% of the GDP and together with agro-based products fetches 80% of the country's total export earnings. More than 43.7% of the labor force is engaged in this sector. [4] The crops are most vibrant sub-sector of agriculture, it consists of 39.6% of agriculture and 8.3% of GDP. Therefore, any change in agricultural productivity creates a ripple effect throughout the rural population of Pakistan. Thus, rapid agricultural growth can stimulate and sustain the pace of industrial growth, setting into motion a mutually reinforcing process of sustained economic growth. [5]

Punjab contributes about 76% to annual food grain production in the country. According to the Punjab Agriculture department "Punjab has 57% of the total cultivated and 69% of the total cropped area of Pakistan. [6] It provides about 83% of cotton, 80% of wheat, 97% fine aromatic rice, 63% of sugarcane and 51% of maize to the national food production. Among fruits, mango accounts for 66%, citrus more than 95%, guava 82% and dates 34% of total national production of these fruits. [7]

1.5 Wheat Production in Pakistan

Wheat flour or "Atta" is the common food for most Pakistanis, supplying 72% of caloric energy in the average diet. It is the main food cereal crop in Pakistan and is essentially better from nutritional point of view than most of the cereals and other foods. It is estimated that in our country wheat consumption per capita is 124 kg per year, which is the highest amount over the world. [5] The above fact reflects the importance of wheat crop for our country. Therefore, every year variation in the total yield of wheat due to several factors including climate and weather variations directly affects the economy and social balance of the country.

Wheat is a Rabi crop that is grown in the winter season. In Pakistan sowing of wheat takes place from October to December and harvests from the month of March to May. In Punjab sowing months of wheat are November and December whereas harvesting period is April and May.

Chapter 2

MATERIALS AND METHOD

The wheat crop variety Arooj-2022 was cultivated at Regional Agrometeorological Center, Faisalabad in central Punjab during Rabi season 2022-23. Both the meteorological and crop phenological data were recorded during the crop season. In order to compile the data of each development stage, careful, precise and timely recording of the following parameters (table 2.1) were undertaken at 0300, 0900 and 1200 (UTC) as routine practice.

Table 2.1: Observed Meteorological Parameters

1.	Air Temperature (°C)
2.	Maximum & Minimum Temperature (°C)
3.	Soil Temperature (°C)
4.	Relative Humidity (%)
5.	Precipitation (mm)
6.	Pan Evaporation at 0300 & 1200UTC
7.	Bright Sunshine Hours
8.	Wind Speed (km/hour)
9.	Wind Direction
10.	Soil Moisture (%)

Crop data including phenological data and soil moisture observation was collected according to World Meteorological Organization (WMO) and Food & Agriculture Organization (FAO) standards.

2.1 Phenology

For a sound understanding of plant growth and development, observation and recording of phenological stages of plant is an essential element (table 2.2). The impact of precipitation, heat, drought, diseases, insects, and weeds can be more accurately predicted with a clear picture of the relationship between growth stage and plant response to stress. The optimum timing of fertilizer, irrigation, herbicide, insecticide and fungicide applications are also best determined by crop growth or phenological stage rather than calendar date. [8]

Table 2.2: Phenological Stages of Wheat Crop 2022-23

	Phenological stage	Date
1.	Sowing	10-11-2022
2.	Emergence	17-11-2022 To 25-11-2022
3.	Third Leaf	26-11-2022 To 04-12-2022
4.	Tillering	05-12-2022 To 08-01-2023
5.	Shooting	09-01-2023 To 12-02-2023
6.	Heading	13-02-2023 To 21-02-2023
7.	Flowering	22-02-2023 To 07-03-2023
8.	Milk Maturity	08-03-2023 To 24-03-2023
9.	Wax Maturity	25-03-2023 To 02-04-2023
10.	Full Maturity	03-04-2023 To 08-04-2023

2.1.1 Phenological Observations

Generally, the field selected for Phenological observations should be of one hectare in size but in this case area of field selected for observation was half acre and it was divided into 4 replications. Over all 10 plants were selected from each replication. These plants were tagged in a row in each replication. Thus, phenological observations were recorded on 40 plants and continued throughout the period on the same plants.

Total number of plants in a particular phenological phase at the same time was observed from each replication on every Monday, Wednesday and Saturday and these observations were recorded on the prescribed Performa. When 10% of the selected plants were in certain phase, that particular phase was considered to be started. If 50% of the selected plants displayed a certain phase, that phase was considered to be in full swing. Similarly, 75% occurrence of certain phase displayed by the selected plants was considered as completion of that particular phase and next Phenological phase observations were started at their proper time. Thus, next phenological stage is not bound to appear after the completion of first one.

It has been observed that at a time two phenological phases or no phase can exist.

2.2 Methodology

In this study the impact of different meteorological/non-meteorological parameters in comparison with the normal (1981-2010), including soil temperature and soil moisture, amount of seed per acre cultivated, time of sowing, fertilizer intake, number and timing of irrigation water added and pest attack/pesticide used over crop growth, development and final yield are discussed.

Chapter 3

RESULTS AND DISCUSSION

In this chapter, variations in meteorological and non-meteorological parameters and their impact on the wheat crop is analyzed along with a brief summary of whole wheat crop's life cycle. Normally the wheat crop takes 120-140 days to get fully mature. The crop is cultivated in the start of November up to the mid of December. In order to investigate the chief causes about variation in the crop's growth, development and yield related to varying weather conditions and some other factors during each phenological stages of wheat crop, different meteorological parameters are studied as well.

Table 3.1: Brief Summary of the Wheat Crop

1	Field size	4 kanal
2	Crop variety	Arooj-2022
3	Date of Sowing	10-11-2022
4	Information about any disease/pest attack,	Aphid, Rust
5	Pesticides And weedicides details	Buctrilsaper 300ml/Hec, Axil 330 ml/Hec
6	Quantity of seed per acre	50 Kg
7	Row spacing	30cm
8	Schedule and quantity of supplied dose of fertilizer	1 bag Potash,2 bag DAP at sowing 1 bag Urea at 1 st Irrigation
9	Type of irrigation	Flood irrigation
10	Irrigation schedule	06-12-2022 (First irrigation) 11-01-2023 (Second irrigation) 24-02-2023 (Third irrigation) 07-03-2023 (Fourth irrigation)
11	Heat units consumed from sowing to full maturity	1695
12	Total days taken by the crop from sowing to full maturity	143
13	Date of harvesting	18-04-2023
14	Actual/ Potential yield	2040/3000 kg /acre

3.1 Rainfall and Wheat Crop Growth

Rainfall is one of the most important factors that affect annual wheat production in Pakistan. In time or effective rainfall i.e., before sowing and shooting to grain formation stage is greatly beneficial. In the same way, rain just after sowing and before germination and at the time of full maturity negatively affect crop growth and ultimately reduce the yield. [9]

During the crop season 2022–23, The months of November, January, March and April received above than normal rainfall while December and February received below rainfall. March received the highest amount of rainfall. During the Month of January, a spell of 30.6mm occurred on 29th January during shooting stage. Overall 62% above Normal precipitation was observed during the crop life.




Figure 3.1: Comparison of Monthly observed rainfall with Normal (1981-2010) at Faisalabad during Rabi season 2022-23

Table 3.2: Daily Rainfall History of the Crop Life 2022-23

Year	Phenological stage	Month	Day	Rainfall(mm)	Total Precipitation During Stage (mm)	Monthly Total (mm)
2022	Sowing	November	10	0.0	0.0	
2022	Germination	November	13	3.0		
2022	Germination	November	14	1.0	4.0	
2022	Emergence	November	17	--	0.0	
2022	Third Leaf	November	26	--	0.0	4.0
2022	Tillering	December	28	0.9		
2022	Tillering	December	29	1.4	2.3	2.3
2023	Shooting	January	11	0.1		
2023	Shooting	January	18	TR		
2023	Shooting	January	19	TR		
2023	Shooting	January	29	30.6		30.7
2023	Shooting	February	05	TR		
2023	Shooting	February	08	TR	30.7	
2023	Heading	February	13	--	0.0	
2023	Flowering	February	27	4.4		
2023	Flowering	February	28	2.0		6.4
2023	Flowering	March	01	TR	6.6	
2023	Milk Maturity	March	16	3.2		
2023	Milk Maturity	March	17	0.6		
2023	Milk Maturity	March	21	TR		
2023	Milk Maturity	March	22	TR		

2023	Milk Maturity	March	23	12.0		
2022	Milk Maturity	March	24	14.4	30.2	
2023	Wax Maturity	March	29	6.0		
2023	Wax Maturity	March	30	13.0		
2023	Wax Maturity	March	31	6.0		55.2
2023	Wax Maturity	April	02	17.4	42.4	
2023	Full Maturity	April	03	6.0		
2023	Full Maturity	April	05	1.4		
2023	Full Maturity	April	17	3.2	10.6	
2023	Harvesting	April	18	--		00

3.2 Irrigation during Crop Growth

The wheat crop was irrigated four times during the entire crop season before full maturity. First irrigation was made to the wheat crop 26 days after sowing during tillering stage. During the shooting stage, second irrigation was made. Third irrigation was made during flowering stage. Fourth irrigation was made at the start of milk maturity stage.

3.3 Air Temperature and Wheat Crop Growth

Air temperature is also one of the most important climatic variables that affect plant life. Plants growth is restricted to certain limits of air temperature. An average increase of 1.9 °C in the mean minimum temperature is recorded in the Punjab over the past 54 years (1960-2013). The increase of 1.4 °C in the mean minimum temperature is projected for the next 30 years. This projected increase is a matter of concern for food and agriculture stakeholders in the Punjab province. [10]

The main dry matter process i.e., photosynthesis is also temperature dependent. Hence three temperature values for a plant growth are of particular importance.

- Biological Zero: is the minimum temperature below which plant growth stops; for wheat crop it ranges between 0°C and 5°C.
 - Optimum Temperature: at which maximum plant growth occurs. For wheat crop its value is 25°C.
 - Maximum Temperature: above which the plant growth stops. For wheat crop its value is 30-32°C.
- [11]

Detail of Mean Monthly Air Temperature, Monthly Mean Maximum and Minimum Temperature and Absolute Maximum and Minimum temperature are presented in Table – 3.3 along with Normal (1981-2010) values and is also shown in following figures (3.2 – 3.4).




Figure 3.2: Mean Monthly Minimum Temperature (°C) of Faisalabad during the Rabi Season




Figure 3.3: Mean Monthly Maximum Temperature (°C) of Faisalabad during the Rabi Season




Figure 3.4: Mean Monthly Temperature during 2022-23

During the crop season of 2022–23, mean day time maximum temperature remained around normal during crop life except the month of February when it was above than normal value. Mean night time minimum temperature also remained slightly above normal during the most of the crop life especially during the month of February and March. Mean daily temperatures also remained around the normal values during most of the crop season expect February when it was above normal.

Table3.3: Mean Monthly Temperatures during Rabi Season 2022-23

Month	Mean Monthly (°C)	Monthly Mean Max.(°C)	Monthly Mean Min(°C)	Absolute Max. (°C)	Absolute Min. (°C)
Nov-2022	20.3 (19.2)	28.2 (27.2)	12.4 (11.5)	33.5 (34.0)	8.0 (2.5)
Dec-2022	14.6 (13.7)	21.8 (21.6)	7.4 (6.3)	27.0 (29.2)	3.0 (-1.3)
Jan-2023	11.6 (11.8)	18.4 (18.7)	4.8 (5.0)	23.0 (26.2)	1.0 (-1.0)
Feb-2023	18.4 (14.8)	26.1 (22.0)	10.7 (7.9)	30.5 (30.8)	6.8 (0.0)
Mar-2023	21.8 (20.0)	28.1 (27.6)	15.4 (13.7)	32.5 (37.0)	12.0 (3.5)
April-2023	26.0 (26.2)	33.6 (34.1)	18.4 (18.9)	39.5 (44.0)	14.5 (7.0)

() in Table 3.3 shows Normal values (1981-2010)

3.4 Soil Moisture Observations during Crop Growth

Soil moisture plays a vital role during crop’s life. Soil moisture content is proportional to rainfall and intake of irrigated water and is inversely proportional to evapotranspiration from the plant and its surroundings. Variation in soil moisture during crop’s life play important role in plant growth and development. Water or soil moisture requirement of wheat crop varies during different growth or Phenological stages. Highest amount is needed during flowering stage followed by grain formation stages and then vegetative stages [12].

To calculate soil moisture, soil samples are taken on 7th, 17thand 27th of each month in four replications at 5, 10, 20, 30, 40, 50, 70 and 90 cm depths with the help of auger. However, in case of any anomalous event on the specific date, the sample can also be taken on the next day. Soil sample is then weighed and dried in the oven for about 8 hours. The dried sample is weighed again and moisture present in the soil is then calculated by the difference of weight between wet and dry samples as illustrated below:

$$\text{Moisture}(\%) = \frac{(\text{Weight of the cane containing soil before drying} - \text{Weight of the cane containing dry soil})}{(\text{Weight of cane containing dry soil} - \text{weight of cane})} \times 100$$

Moisture contents of the soil varied due to dry and wet spells throughout the season. After each effective irrigation or rain the moisture level increased in the shallow layers as compared to deep layers of the soil.

During the crop season 2022–23, from the observed soil moisture data, figures (3.5 – 3.7) depicts that soil moisture remained satisfactory in general. It remained deficit at lesser depths during the early growth stage. The temperature remains normal during this period. Rainfall received during the crop season was above normal. Vertical axis shows the depth (cm) at which soil sample is taken and horizontal axis shows the date of soil sample collection.




Figure 3.5: Soil moisture chrono Isopleths for Rabi Crop at Faisalabad for the year 2022-23.




Figure 3.6: Soil moisture at different depths during wheat crop 2022-23



Figure 3.7: Soil moisture at deep soils (90cm) during wheat crop 2022-23

3.5 Soil Temperature and Crop Growth

Soil temperature is the most important parameter affecting the growth of the crop. Plants roots are very sensitive to the soil temperature. In comparison to air temperature, the amplitude of variation in soil temperature is much more pronounced because of the varying characteristics, texture, composition, and organic material of soil. Soil temperature influences the germination of seeds, the functional activity of the root system, the incidence of plant diseases and the rate of plant growth. The daytime soil temperature is more important than the night time temperature, because it is necessary to maintain the internal crop water status to match the evaporation rate. Optimal soil temperature for growth of wheat plant roots during the vegetative stage is below 20°C and is further lower than that for the shoots. [11]

Soil temperature is also an easy tool to predict the status of soil moisture content during varying conditions of air and soil throughout crop's life. Generally, above normal condition of soil temperature at a particular depth indicates deficiency of soil moisture content and below normal soil temperature indicates satisfactory condition of soil moisture content.

From the observed data, it is evident that generally, soil temperature increases gradually with increasing depths. Soil temperature varies as soil moisture varies from depth to depth and time to time. To measure soil temperature, the soil thermometers were installed at different depths to monitor the thermal regime of the soil. The soil temperatures in degree Celsius (°C) were observed and then recorded three times a day at 0300, 0900 and 1200 UTC. The depth at which the soil temperatures were observed on daily basis includes 5, 10, 20, 50 and 100 cm depths. It was observed that major root concentration is up to 40 cm.

Note: This soil temperature data is collected from the soil observations taken at agromet observatory of RAMC Faisalabad situated near the experimental field of wheat crop. Therefore, this data tells us a general status of soil moisture of the soils of the area (which is not irrigated) and not of the crop's field particularly, which is irrigated as per requirement. It is thus important to note that any deficiency in soil moisture indicated by soil temperature data may or may not be actually experienced by the crop's soil, which was irrigated in accordance with water requirement of the crop several times during its life time.

During the crop season 2022-23, soil temperature remained highest at all depths during month of April except for 100cm depth which was recorded highest during the month of November. It remained coolest in the month of January at all depths. (Data for 50cm depth is not available)




Figure 3.8: Soil temperature during wheat crop 2022-23

3.6 Heat Units Consumption during Crop Cycle

Heat units, Growing Degree Days, effective heat units or growth units are a simple means of relating plant growth, development, and maturity to air temperature. Heat units are often used in agronomy, essentially

to estimate or predict the length of the different phases of development in plants.

The heat unit concept assumes a direct and linear relationship between plant growth and temperature. It starts with the assumption that total plant growth is dependent on the total amount of heat to which it is subjected during its life time. The heat units for a particular crop on any day are the difference between the daily mean temperature (T) and the base temperature (T_b) of the crop. Base temperature or Biological zero is the minimum temperature below which no growth occurs. The base temperature or threshold varies with different plants, and for the majority it ranges from 4.5 to 12.5°C. Here for wheat crop base temperature is 5°C. This concept assumes that a given cultivar requires the same summation (k) of the daily mean temperature for growing from one phenological stage to next stage, regardless of temperature distribution. Generally, only positive values above the biological zero (T_b) are considered. The period of negative value is termed as “dormant” because crop does not grow under such conditions.

In this study the effective method has been used for determining heat units which is represented by the following equation.

$$GDD = \sum(T - T_b) = k \quad \text{if } T > T_b \text{ and } GDD = 0 = k \quad \text{if } T < T_b$$

Where *T* = Mean daily temperature, *T_b* = Biological Zero (5°C), *k* = Heat Unit [10].

Inter Phase period for wheat crop during 2022-23 and corresponding heat units at RAMC Faisalabad observed at different phenological stages varies from phase to phase. Heat unit requirements of different phases and cumulative heat units for the crop have been worked out, are shown in figure 3.9 and Table-3.4. Total heat units consumed by the wheat crop were 1694.6 accumulated from germination to full maturity in 143 days. On average 11.8 heat units were consumed by the crop per day. Normally these heat units are consumed in 140 days at the rate of 11.6 heat units per day, which shows that the crop reached to full maturity in almost normal time. From figure 3.9 it is clear that crop consumed normal to above normal heat units during most of the vegetative stages. Below normal heat units were consumed during reproductive stages of flowering and full maturity. Milk Maturity stage again consumed higher than normal value of heat units. So most of the crop period experienced normal to higher than normal heat units.




Figure 3.9: Heat units during crop life 2022-23

Table 3.4: Heat Units Consumed by the Crop during Different Phenological Phases

S. No.	Inter Phase	Period	No. of Days Taken	Degree Days (T-5°C)	Normal Degree Days
2.	Emergence	17-11-2022 To 25-11-2022	09	115.5	132.5
3.	Third leaf	26-11-2022 To 04-12-2022	09	110.5	111
4.	Tillering	05-12-2022 To 08-01-2023	35	290.9	241
5.	Shooting	09-01-2023 To 12-02-2023	35	300.8	237
6.	Heading	13-02-2023 To 21-02-2023	09	125.3	131
7.	Flowering	22-02-2023 To 07-03-2023	14	225.7	286
8.	Milk maturity	08-03-2023 To 24-03-2023	17	293.8	257
9.	Wax maturity	25-03-2023 To 02-04-2023	09	127.4	122
10.	Full maturity	03-04-2023 To 08-04-2023	06	104.7	154
11	Emergence to Maturity	17-11-2022 To 08-04-2023	143	1694.6	1672

3.7 Relative Humidity (%)

Relative humidity (RH) is simply defined as the ratio of the actual amount of water vapor (grams) in the unit mass of air at a given temperature to the mass of water vapor in the same sample of air when it is saturated at the same temperature. RH tells us about how much a sample of air around the observing station is saturated. It is expressed in percentage. For example, if relative humidity of the atmosphere is 60%. It means that 60% of the atmosphere is saturated with water vapor or moisture and 40% of the present moisture may be added more to saturate (100% RH) the atmosphere. RH is temperature and moisture dependent and may vary if moisture content or temperature of the atmosphere is changed. It is always inversely proportional to temperature and ETo. RH is calculated with the help of dry bulb and wet bulb temperatures fitted in the Stevenson screen. During the crop season 2022-23, RH remained below than the normal values during the vegetative growth period but it remained normal during the reproductive growth period.

3.8 Reference Crop Evapotranspiration, ETo (mm/day)

Reference Crop Evapotranspiration or ETo is defined as the rate of Evapotranspiration from an extended surface of 8– 15 cm tall green grass cover of uniform height, actively growing, completely shading the ground, free from disease and not short of water. The standard method of calculating ETo is revised Penman-Monteith equation. Using this method, ETo is calculated using data of temperature (maximum and minimum), wind speed, relative humidity and solar radiation. Solar radiations are calculated with the help of total bright sunshine hour data at a particular station. Sunshine duration recorders with sunshine cards are used for sunshine hours recording [12].

During the crop season 2022-23, relative humidity remained below normal during the vegetative growth period but it remained normal during the reproductive growth period whereas, ETo also remained below normal during the whole crop life.

3.9 Wind and Crop Growth

Wind also play significant role in plant growth besides its role in variation of ETo. Normal/gentle wind is necessary for the movement of carbon dioxide to plant canopy so that normal rate of photosynthesis continue in day time. Strong cyclonic or stormy wind accompanied by any severe weather event like hail storm, heavy shower may badly affect/damage the crop. During the Rabi crop period 2022-23, a few such events occurred during the crop life but wind gust recorded was below 15 knots in each of the cases and no damaged was reported. Average wind speed remained below normal during most of the crop life.

Table 3.5: Summary of some Meteorological Parameters during Rabi Season 2022-23

Month	Wind speed (km/hr)	RH (%)	Days with mean RH \geq80%	ETo (mm/day)
Nov-22	1.6 (1.0)	57 (71.2)	00	2.1
Dec-22	1.4 (1.4)	65(74.7)	05	1.4
Jan-23	1.7 (2.3)	63 (76.5)	05	1.2
Feb-23	1.8 (3.0)	53 (71.2)	00	2.1
Mar-23	2.4 (3.9)	64 (65.6)	03	2.9
Apr-23	2.7 (3.6)	50 (50.8)	00	4.4

2.1. Crop Water Requirement (CWR)

The crop water requirement (ET crop) is defined as the depth (or amount) of water needed to meet the water loss through evapotranspiration. In other words, it is the amount of water needed by the various crops to grow optimally.

The crop water need always refers to a crop grown under optimal conditions, i.e. a uniform crop, actively growing, completely shading the ground, free of diseases, and favorable soil conditions (including fertility and water). The crop thus reaches its full production potential under the given environment.

The crop water need mainly depends on:

The climate: in a sunny and hot climate crops need more water per day than in a cloudy and cool climate.

The crop type: crops like maize or sugarcane need more water than crops like millet or sorghum.

The growth stage of the crop; fully grown crops need more water than crops that have just been planted [12]. Sajjad et al., used the output dataset from the Coordinated Regional Climate Downscaling Experiment (CORDEX) for South Asia, under two emission scenarios (RCP 4.5 and RCP8.5) concluded that future CWR is continuously increasing over the entire region, whereas significant CWR increase was observed for the months of August to November over northern and southern parts during the near future (2011-2040). The highest significant increase in CWR rate was recorded in August (0.24 mm/day) and September (0.22 mm/day) over Pakistan at 95% confidence level. The highest significant increase in CWR rate was observed in August (0.56 mm/day) and March (0.30 mm/day) over the whole country. Projected seasonal CWR indicated significant increase under RCP4.5 for Rabi seasons (0.43 mm/day) over the entire country during near future (2011-2040) [14].

3.10 Calculation of Crop Water Requirement (CWR)

After determining E_{To} , the ET crop/crop water requirement (CWR) can be predicted using the appropriate crop-coefficient (K_c)

$$ET_{crop} = K_c \cdot E_{To} \quad \text{or} \quad CWR = K_c \cdot E_{To}$$

Crop coefficient (K_c) is actually the ratio of maximum crop evapotranspiration to reference crop evapotranspiration. For wheat, this ratio becomes 1 during the reproductive cycle (heading to grain formation) otherwise it remains less than 1 bearing minimum values during the early age of the crop and at maturity. The crop water requirement was calculated for the period from emergence to wax maturity. After wax ripeness practically, there is no need of irrigation because the hot and dry conditions are desirable to achieve rapid hard maturity. A schematic variation of the crop coefficient related to different crop development stages under normal conditions is given in figure 3.10.




Figure 3.10: March of Crop Coefficient (K_c) for normal duration of Wheat growing season (Emergence to Wax- Maturity).

The water requirements can be calculated in millimeters and they can be converted into cubic meter per hectare by following equation:

$$10 \text{ mm} = 1 \text{ cubic meter per hectare}$$

Along with the loss of water through evaporation and transpiration, the compensation of this loss by precipitation may also be considered. Normally most of the plants grow successfully and utilize water for

the soil at 50% and above available soil moisture. The maximum demand (daily or seasonal) may be equal to the reference crop evapotranspiration (ET_o) which is utilized through soil moisture [15].

During the Rabi Season 2022-23 in Faisalabad, crop water requirement of wheat crop was observed below normal during the crop life. (Fig 3.11& Table 3.6). ET_o also are remained below normal during all of the phenological stages. The available irrigated and rain water satisfied moisture requirement of the crop, which resulted in normal crop growth.




Figure 3.11: Crop Water Requirement (CWR) During Crop Life

Table 3.6: Heat Crop water requirement during different phenological phases

S. No.	Inter Phase	Period	No. of Days Taken	ET _o (mm)	*ET _o (mm)	CWR=Kc ET _o	CWR=Kc *ET _o
2.	Emergence	17-11-2022 To 25-11-2022	09	19.4	29.5	13.15	20
3.	Third leaf	26-11-2022 To 04-12-2022	09	17.1	24	12.97	18.2
4.	Tillering	05-12-2022 To 08-01-2023	35	43.8	74	40.37	68.2
5.	Shooting	09-01-2023 To 12-02-2023	35	48.3	64	44.22	58.6
6.	Heading	13-02-2023 To 21-02-2023	09	17.6	32.5	12.24	22.6
7.	Flowering	22-02-2023 To 07-03-2023	14	32.2	74.1	16.08	37
8.	Milk maturity	08-03-2023 To 24-03-2023	17	52.1	62.1	16.28	19.4
9.	Wax maturity	25-03-2023 To 02-04-2023	09	27.6	32.4	5.54	6.5
10.	Full maturity	03-04-2023 To 08-04-2023	06	23.8	37.8	3.59	5.7
11	Emergence- Full maturity	17-11-2023 To 08-04-2023	143	281.9	430.4	164.4	256.2

*Normals based upon 1991-2010 data.

3.10 Agro Meteorological Summary of Crop Cycle

Different meteorological parameters were recorded at various phases of wheat crop during rabi season 2022-23. The impact of these parameters at different phenological stages of wheat crop is discussed as under.

3.10.1 Sowing

Wheat crop was sown at the start of second decade of November which is recommended/normal time for sowing. The mean temperature was around normal value and there was absolutely no rain after sowing so overall conditions were satisfactory for sowing.

3.10.2 Emergence

Emergence phase was distinguished by the appearance of spike above the ground. When plant emergence stage was completed; the field was divided into four replications. The mean relative humidity during this phase was 51%. Mean air temperature during emergence was 18.4°C. No rainfall was reported during this stage. Crop growth was reported satisfactory and no abnormal weather was reported during emergence.

3.10.3 Third Leaf

This phase is characterized by the appearance and extension of the third leaf to 1-2 centimeter length. This phase took 05 days to complete. The mean relative humidity was 60%; mean air temperature was 17.2°C. This phase experienced no rainfall. Normal crop growth was reported during this stage.

3.10.4 Tillering

This stage of crop growth took 06 days to complete. The mean air temperature during this phenological stage was 13.3°C and mean relative humidity was 67%. Rainfall amount recorded was 2.3 mm during this stage.

3.10.5 Shooting

During this phase stem extension occurred in the crop. The mean air temperature was 13.6°C and mean relative humidity was 56% during this stage. Rainfall received was 30.7mm during this phase.

3.10.6 Heading

It is the initiation of reproductive stage of wheat crop. The mean air temperature was 19.2°C during the heading phase. The relative humidity was around 52%. No rain was received during this stage.

3.10.7 Flowering

The mean air temperature during this stage was recorded as 21.1°C and mean relative humidity was 60%. Rainfall reported was 6.6 mm during flowering.

3.10.8 Milk Maturity

Seed formation is considered as the most important phenological stage of the plant growth. The mean air temperature was recorded as 22.3°C whereas mean relative humidity during this phase was 65%. Precipitation of 30.2 was recorded during this phase. This stage took 08 days for completion.

3.10.9 Wax Maturity

The mean air temperature was 21.3°C and relative humidity was recorded as 65% during wax maturity phase. This stage took 08 days for its completion. This phase experienced 42.4 mm of rainfall.

3.10.10 Full Maturity

Wheat requires high temperature at this stage for maturity. This phase took 06 days to complete. The mean air temperature recorded during this phase was 22.4°C and the relative humidity was 52%. Rainfall observed was 10.6mm during this phase.

Chapter 4

CONCLUSION AND RECOMMENDATIONS

The crop variety Arooj-2022 was cultivated in the experimental field of Ayub Agriculture Research Institute Faisalabad. The sowing time of wheat is a very important factor. The time of sowing of wheat is directly related to yield and quality of grain produced.

According to Table – 3.1 and Chapter-3 (Results and Discussion), the crop was sown at the proper time. Amount of seed cultivated per acre was 50 kg which is sufficient for the crop cultivated up to the start of December. Fertilizer intake was also enough i.e., 2 bags DAP and 1 bag Potash at the time of sowing were added to the crop while 1 bag urea during 1st irrigation was added. Above normal rainfall was reported in the month of January while overall rainfall was much below normal during the crop life. Irrigation was made three times during the crop season which mostly fulfilled the water requirement of the crop. Day time temperatures were observed mostly in normal range. Night temperatures were observed mostly above normal values. RH was observed to be below normal to normal during the crop season. However, ETo remained below normal during the crop's life. As a result, normal moisture content of soil in major root zone was observed during most of the crop life thus contributing to normal crop's growth and development during vegetative stages. Total heat units consumed by the wheat crop were 1695 accumulated from emergence to full maturity in 143 days. On average 11.8 heat units were consumed by the crop per day. Normally these heat units are consumed at the rate of 11.6 heat units per day. From figure 3.9 it is clear that crop consumed above normal heat units at tillering, shooting and milk maturity stages for all the remaining stage it used below normal heat units.

Excess of weeds also play important role in yield's reduction as weeds consume considerable amount of moisture and other soil nutrients and negatively affect crop's growth at the same time. But this issue can be resolved by proper and timely use of recommended varieties of weedicides. During 2022-23 recommended weedicides operations were performed in time, which also reinforced crop's growth and production. Widely spread Rust attack was observed on the crop during its life cycle.

4.1 Conclusions

During the Rabi season 2022-23, most of the air and soil weather parameters like soil temperature, rainfall, irrigation with suitable intervals, R.H and crop water requirement were in range required for normal crop growth. The normal day time and mean temperatures favor the normal completion of reproductive stages of grain formation and development. The favorable weather conditions resulted in high yield of the crop.

4.2 Recommendations

Keeping above results and conclusions, following recommendations/suggestions are given to farmers and other stake holders to enhance wheat crop yield in central Punjab as well as all over Pakistan.

1 – Farmers generally plant wheat late in rice-wheat, sugarcane-wheat, and cotton-wheat areas due to late harvesting of Kharif crop which results in drastic low yields because the crop is exposed to heat stress at grain formation stages (milk and wax maturity stages) leading to the formation of shriveled grain. Late-planted crop has lower germination, fewer tillers, smaller heads, shriveled grain and lower biomass than the timely planted crop. Any delay in planting would reduce yield drastically. To achieve good yield, sowing of wheat should be completed up to the mid of November in the irrigated planes of Punjab.

2 – Wheat plant water requirement is maximum during flowering and grain formation stages (milk and wax maturity stages) followed by vegetative stages. Therefore, farmers and other decision makers should make possible the availability of irrigation water to wheat crop keeping this order in mind to get maximum crop yield.

3 – Diseases, especially rusts (Stripe/Yellow Rust, Stem Rust, and Leaf Rust) and emerging scenario of increased incidences of Powdery Mildew and aphid are major biotic stresses of wheat crop that inflict heavy losses when in epidemic form. A simple one-year disease epidemic could cause a 2-3 billion-rupees loss to the country because of reduction in wheat yield (an example of 1977-78 epidemics). Therefore, breeders and pathologists should join hands to develop disease resistant and high yielding varieties in order to cope with threats created by ever changing rust races by using approaches like durable resistance. Farmers should use approved varieties of pesticides and insecticides.

4 – The frequency of extreme weather events like heat waves, cold waves, flash flooding, and heavy snowfall has increased globally including Pakistan in the last decade due to climate change. Pakistan has also been facing water shortages and drought conditions for the last several years due to lesser rains and high temperatures due to global warming which resulted in hampering of wheat production both in irrigated and rainfed areas. In order to minimize the negative effects of climate change and accompanied global warming, drought and heat tolerant varieties need to be evolved in addition to the judicious use of available irrigation water. Researches should also keep focus on to update sowing dates due to late arrival of winter season and winter rainfall in most of the agricultural plains of Pakistan. Keeping in mind the available water resources, it is also indeed necessary to decide suitable crops to be cultivated in a particular region.

5 – Farmers may be advised to be in contact with local and Federal Agricultural Departments and Pakistan Meteorological Department throughout crop's life, especially at the time of sowing, adding fertilizers to crop, before irrigation and harvesting. It will help the farmers to get in time weather advices to deal in

better way with any present or coming water stress condition and to get best results of fertilizer and irrigated water used.

6 – Weeds being the main robbers of plant food from soil; space and even light required for wheat plants, be controlled by cultural practices and in case of heavy infestation, may be eliminated by application of recommended herbicides and weedicides. This technique will definitely increase the yield.

7 – Seed of high yielding wheat varieties resistant to rusts, smuts, etc. approved by Agricultural Department for a particular region in a particular amount must be used. Seed should be treated with a suitable insecticide carefully before sowing.

8 – Timely application of nitrogen-phosphoric fertilizers should be done.

9 – Care must be taken to check the pre and post-harvest losses of wheat. Pre-harvest damage may be checked from attack of birds, animals around fields. Post-harvest losses must be checked from the attack of rodents and other insects, pests and fungi. After proper trashing, wheat grains should be placed under hygienic conditions in fields and in storage places as well.

10 – Crop rotation is an important factor that enriches the fertility of the land, which should not be ignored. Pulses are preferred as they enrich the nitrogen content of soil for the coming wheat crop.

References

- [1] Chaudhry, Q. Z. and G. Rasul (2004): Agroclimatic Classification of Pakistan, Science Vision Vol.9 No.1-2 (Jul - Dec, 2003) & 3-4 (Jan - Jun, 2004).
- [2] Shamshad, K.M. (1988): Meteorology of Pakistan, Royal Book Agency, Karachi, Pakistan.
- [3] Monthly Climatic Normals of Pakistan, 1981-2010 (January 2013): Climate Data Processing Center, Pakistan meteorological Department, Karachi.
- [4] Economic Survey of Pakistan, 2013-14: <http://budget.par.com.pk/dll/1189/>
- [5] Punjab Agriculture Department 2014: <http://www.agripunjab.gov.pk>
- [6] Pakistan Agricultural Department: <http://www.defence.pk/forums/national-political-issues/102679-punjab-door-revolution.html#ixzz2PWvbxY6r/>

- [7] USDA Foreign Agriculture Service Report (2014): www.fas.usda.gov
- [8] Fowler, D. B. (2002): Growth stages of Wheat, Chapter 10: http://www.usask.ca/agriculture/plantsci/winter_cereals/index.php
- [9] Imran, A, Q. Zaman and M. Afzal (2013): Temporal Trends in the peak monsoonal precipitation events over Northeast Pakistan, Pakistan Journal of Meteorology. Vol. 10, Issue 19: July 2013.
- [10] Sajjad, H and A. Ghaffar (2019): Observed, simulated and projected extreme climate indices over Pakistan in changing climate, Theoretical and Applied Climatology (2019) 137:255–281 <https://doi.org/10.1007/s00704-018-2573-7>
- [11] Mavi, H. S. and Graeme J. T. (2005): Agrometeorology; principals and applications of climate studies in agriculture, international book distributing company, Lucknow, India.
- [12] FAO Irrigation and Drainage Paper 33.
- [13] Crop water needs chapter10. FAO. (<http://www.fao.org/docrep/S2022E/s2022e07.htm>).
- [14] Sajjad, H and K. Ullah (2019): Projected crop water requirement over agro-climatically diversified region of Pakistan, Agricultural and Forest Meteorology, www.elsevier.com/locate/agrformet
- [15] Naheed, G. Rasul (2009): Water requirement of Wheat crop in Pakistan, an MS dissertation, COMSATS, Islamabad.